

### Elementary Dynamics Example #3: (Rectilinear Motion)

Given:  $a(v) = g - cv$  ( $a(v) \geq 0$ ) ... the **acceleration** of an object falling in a fluid.

Initial condition:  $v(0) = 0$

Find:  $v(t)$  ... the **velocity** of the particle as a **function of time**

Solution:

$$a(v) = \frac{dv}{dt} = g - cv \Rightarrow \frac{dv}{g - cv} = dt \Rightarrow \frac{-1}{c} \int \frac{-c \, dv}{g - cv} = \int dt \dots \text{using indefinite integrals}$$

$$\boxed{\frac{-1}{c} \ln(g - cv) = t + D}$$

Applying the **initial condition**,  $v(0) = 0$ , gives  $\boxed{D = \frac{-1}{c} \ln(g)}$

Aside:

$$\int \frac{f'(x) \, dx}{f(x)} = \ln(f(x))$$

**Substituting** the value of the constant  $D$  into the first boxed equation and **simplifying**:

$$\frac{-1}{c} \ln(g - cv) = t + \frac{-1}{c} \ln(g) \Rightarrow \ln(g - cv) = \ln(g) - ct \Rightarrow g - cv = \exp(\ln(g) - ct)$$

$$g - cv = \exp(\ln(g)) \cdot \exp(-ct) = g e^{-ct} \Rightarrow \boxed{v(t) = \frac{g}{c} (1 - e^{-ct})}$$

The function  $v(t)$  starts at **zero** and increases exponentially to the **final value** of  $\frac{g}{c}$ . The **larger** the value of the coefficient  $c$ , the **faster** the function increases toward  $\frac{g}{c}$ . The plot below shows a plot of  $v(t)$  for  $c = g = 1$ .

